

Research

Open Access

An encoded N-terminal extension results in low levels of heterologous protein production in *Escherichia coli*

Samantha S Orchard² and Heidi Goodrich-Blair*¹

Address: ¹Department of Bacteriology, University of Wisconsin, Madison, WI 53706, USA and ²Department of Biology, San Diego State University, San Diego, CA 92182 USA

Email: Samantha S Orchard - samantha_orchard@yahoo.com; Heidi Goodrich-Blair* - hgblair@bact.wisc.edu

* Corresponding author

Published: 21 July 2005

Received: 14 June 2005

Microbial Cell Factories 2005, 4:22 doi:10.1186/1475-2859-4-22

Accepted: 21 July 2005

This article is available from: <http://www.microbialcellfactories.com/content/4/1/22>

© 2005 Orchard and Goodrich-Blair; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The *tdk* gene (encoding deoxythymidine kinase) of the gamma-proteobacterium *Xenorhabdus nematophila* has two potential translation start sites. The promoter-distal start site was predicted to be functional based on amino acid sequence alignment with closely related Tdk proteins. However, to experimentally determine if either of the two possible start codons allows production of a functional Tdk, we expressed the "long-form" (using the promoter-proximal start codon) and "short-form" (using the promoter-distal start codon) *X. nematophila tdk* genes from the T7 promoter of the pET-28a(+) vector. We assessed Tdk production and activity using a functional assay in an *Escherichia coli tdk* mutant, which, since it lacks functional Tdk, is able to grow in 5-fluorodeoxyuridine (FUdR)-containing medium.

Results: Short-form Tdk complemented the *E. coli tdk* mutant strain, resulting in FUdR sensitivity of the strain. However, the *E. coli tdk* mutant expressing the long form of *tdk* remained FUdR resistant, indicating it did not have a functional deoxythymidine kinase enzyme. We report that long-form Tdk is at least 13-fold less abundant than short-form Tdk, the limited protein produced was as stable as short-form Tdk and the long-form transcript was 1.7-fold less abundant than short-form transcript. Additionally, we report that the long-form extension was sufficient to decrease heterologous production of a different *X. nematophila* protein, NilC.

Conclusion: We conclude that the difference in the FUdR growth phenotype between the *E. coli tdk* mutant carrying the long-or short-form *X. nematophila tdk* is due to a difference in Tdk levels. The lower long-form protein level does not result from protein instability, but instead from reduced transcript levels possibly combined with reduced translation efficiency. Because the observed effect of the encoded N-terminal extension is not specific to Tdk production and can be overcome with induction of gene expression, these results may have particular relevance to researchers attempting to limit production of toxic proteins under non-inducing conditions.

Background

Proteins from one organism are often expressed in a different species for the purpose of protein purification or complementation studies. When such efforts fail due to

non-production of the protein, the underlying cause of failure is often unclear [1]. Protein overproduction is known to induce a heat shock-like response, which results in increased proteolysis in the cell and therefore possible





Figure 2

Immunoblot analysis of protein production using anti-Tdk serum (panels A and C-G) or anti-NilC serum (panel B). *E. coli* KY895 (panels A, B, F, G), BL21 Star™ (DE3) (panel C), KY895 *clpP* (panel D), or BL21 (DE3) (panel E) carrying plasmids pTara and pET-28a(+) (vector-only; lane 1), pETSTdk or pETSNiC (short-form; lane 2), pETLTdk or pETLNilC (long-form; lane 3), pETMAGPTdk or pETMAGPNiC (D2A change; lane 4), pETMDAPTdk or pETMDAPNiC (G3A change; lane 5) or pETMDGATdk or pETMDGANiC (P4A change; lane 6) was grown overnight in LB/kan/cam and separated cellular proteins were subjected to immunoblot analysis with the indicated antiserum. Arabinose (0.2%) was added to the culture medium for the strains in panels F and G, and IPTG (0.2 mM) was added to the culture medium for the strains in panel G. Similar volumes of cultures were used in each lane but cultures were not normalized for optical density or overall protein concentrations.

Long-form NilC-trunc is not detected by immunoblot analysis

To determine if the 5' extension of long-form *tdk* would reduce production of a different protein, the 12-nt exten-

sion was added to an *X. nematophila* gene, *nilC*, encoding an outer membrane-associated protein [5] and for which there were readily available antibodies. To create a cytoplasmic version of NilC, the DNA encoding the 21 amino

**Figure 3**

Primer extension analysis of short- and long-form *X. nematophila tdk* transcripts in *E. coli*. Total RNA was isolated from *E. coli* KY895 carrying plasmids pTara and pETSTdk or pETLTdk grown overnight to $OD_{600} = 2.0$. Primer tdkprext2 was end-labeled with ^{32}P and used to produce a sequencing ladder from plasmids pETSTdk (left side; lanes C, T, A, G) and pETLTdk (right side; lanes C, T, A, G). A primer extension product was produced with end-labeled tdkprext2, total RNA template (left side: KY895 carrying pETSTdk; right side: KY895 carrying pETLTdk) and AMV-RT enzyme at $37^{\circ}C$ (lanes "pe"). Arrows indicate the expected transcriptional start site from the T7 promoter of pET-28a(+).

acid long NilC N-terminal signal sequence was removed creating NilC-trunc (short form), to which the MDGP extension was added (long form). Long-form NilC-trunc was not detected by immunoblot analysis with polyclonal anti-NilC serum while short-form NilC-trunc was (Fig. 2B). Thus, the MDGP-encoding extension is sufficient to reduce production of at least two unrelated proteins.

Long-form *tdk* transcript is less abundant than short-form transcript but shares the same 5' end

Primer extension analysis was used to determine if the MDGP-encoding sequence affects the transcription (either level or start site) of *tdk* from pET-28a(+) or processing of the *tdk* transcript. Primer extension analysis indicated that long-form *tdk* transcript was made and at least some portion maintained its 5' end (Fig. 3). Relative transcript levels were quantified by real-time PCR with primers specific for a central portion of the *tdk* gene. For growth in LB broth to $OD_{600} \sim 0.85$, the long-form transcript was 1.7-fold less abundant than the short-form transcript. This lower level of long-form *tdk* transcript could be due to higher levels of long-form *tdk* mRNA degradation,

reduced transcription of long-versus short-form *tdk* or to differences in plasmid copy number, possibly a result of the presence of long-form *tdk* or long-form Tdk production. To determine if this lower transcript abundance is due to increased susceptibility to degradation, protein production was monitored in an *E. coli* RNaseE mutant. RNaseE is one of the main *E. coli* mRNA-degrading enzymes [6] and we hypothesized that it differentially degrades long-versus short-form *tdk* transcript. In the RNaseE mutant, long-form Tdk was detected at low levels by immunoblot analysis (Fig. 2C). Thus in the absence of RNaseE, long-form *tdk* transcript was sufficiently produced and stable to act as a template for protein synthesis, indicating that RNaseE degrades long-form *tdk* transcript. However, even in the absence of RNaseE, long-form Tdk protein levels were ~ 4.7 -fold lower than those of short-form Tdk produced under the same conditions. Therefore, RNaseE-mediated degradation of long-form *tdk* transcript is not the only factor contributing to the low levels of long-form Tdk protein production relative to short-form Tdk.




Figure 4

Quantification of Tdk protein levels and stability. Levels of protein were determined by ^{35}S -methionine labeling of whole cells and subsequent immunoprecipitation with polyclonal anti-Tdk antiserum. *E. coli* KY895 carrying plasmids pTara and pETSTdk (Short-form Tdk) or pETLTdk (Long-form Tdk) was labeled with ^{35}S -methionine for 3 min followed by a chase with excess unlabeled methionine. Samples were removed for processing at 20 s, 2 min and 5 min following the chase and proteins reactive with anti-Tdk serum were immunoprecipitated and separated as described in Methods. Following autoradiography, intensity of the bands corresponding to labeled Tdk were determined and background corrected against a pET-28a(+) vector-only control. The calculated band intensities are expressed in arbitrary units.

Long-form Tdk protein is stable

The stabilities of short- and long-form Tdk proteins were measured to determine if differential susceptibility to post-translational proteolysis is the cause of the low protein levels of long-versus short-form Tdk. Measurement of long-form Tdk stability was made possible by inducing *tdk* expression by addition of arabinose to the culture medium (arabinose induces expression of pTara-encoded T7 RNA polymerase, which is responsible for transcription from the pET-28a(+) promoter controlling *tdk* expression). Pulse-chase labeling of cells and immunoprecipitation experiments revealed that long-form Tdk protein is present at a low level with such induction (Fig. 4), although this level was still 13-fold lower on average than that of short-form Tdk and was insufficient for detection by immunoblotting (Fig. 2F). Over the 5 min time course of the experiment, long-form Tdk was at least as stable as short-form Tdk (Fig. 4), indicating that the difference in protein levels between these two forms is not due to differential proteolysis. Further support for this hypothesis comes from the fact that long-form Tdk was not

detected in *E. coli* KY895 *clpP* and *E. coli* BL21(DE3) (*lon*) protease mutant strains (Fig. 2D–E).

Long-form Tdk is expressed in the presence of IPTG and arabinose

Arabinose was used to induce T7 RNA polymerase for the ^{35}S -methionine labeling study, resulting in detectable long-form Tdk protein levels by autoradiography (Fig. 4) but not immunoblot analysis (Fig. 2F). Neither arabinose nor IPTG (de-repression of the pET-28a(+) T7 promoter) were used for the other protein production studies reported herein. When both IPTG and arabinose were added to cultures of KY895 carrying pTara and pETLTdk, long-form Tdk was detected by immunoblot analysis at levels similar to short-form Tdk in comparably induced cultures (Fig. 2G). Thus, increased transcription of long-form *tdk* is sufficient to promote high long-form protein levels. These data support the hypothesis that the low level of protein expressed from genes with the MDGP-encoding extension results at least in part from a reduction in mRNA levels. These results further indicate that, in non-inducing conditions in cases when overproduction of a toxic protein is desired, the encoded N-terminal extension described here might be beneficial in reducing "leaky" protein production.

Changing the nucleotide and/or amino acid identity of the long-form extension results in variable protein production

To determine the relative importance of each codon in the extension region of long-form *tdk* and *nilC*-trunc to the resulting observed protein levels, we systematically changed the aspartic acid, glycine and proline codons to alanine codons and monitored the resulting effects on protein production. The levels of the altered long-form proteins were assessed by immunoblot analysis following normalization of samples by optical density (Table 2; see also Fig. 2A–B). The only change that caused an increase in the level of both long-form Tdk and *NilC*-trunc protein production was a glycine (GGG) to alanine (GCG) codon change (Fig. 2A and 2B). An aspartic acid (GAC) to alanine codon (GCC) change did not alter the level of production of long-form *NilC*-trunc but did allow detection of long-form Tdk (Fig. 2A). Additional changes were made in the long-form extension region to assess the importance of the position and order of the amino acid residues, particularly glycine (Table 2). All extensions that result in low levels of protein production (similar to that of MDGP) include a glycine codon in the third or fourth position, even when the initiating methionine residue is predicted to be cleaved from synthesized proteins (see Table 2 for a list of extensions where the N-terminal *f*-Met residue is expected to be cleaved by PepM following protein synthesis [7]). Thus, the position of the glycine residue or codon nucleotides appears to impact long-form protein production.

Table 1: Oligonucleotide primers used in this study^a

Oligonucleotide name	Sequence (5'->3') ^b	Target ^c
<u>Bam</u> HI <i>ni</i> C	<u>GGATCC</u> GACGATGCCTTAATGCGACAG	<i>ni</i> C
<u>Bam</u> HI <i>tdk</i>	<u>GGATCC</u> CTTAGTGATTTTATACG	3' of <i>tdk</i>
<i>clpP</i> tetKWfor ^d	ATCGGTACAGCAGGTTTTTTCAATTTTATCCAGGACCG CAAGAGGGTCATTATATTTCCG	5' of <i>E. coli clpP/Tn10</i> (tet ^r)
<i>clpP</i> tetKWrev ^d	GCCGCCCTGGATAAGTATAGCGGCACAGTTGGCCCTCTGG GACTCGACATCTTGGTTACCG	3' of <i>E. coli clpP/ Tn10</i> (tet ^r)
Ecrecaminfor	GAAAGCGAAATCGAAGGCG	<i>E. coli recA</i>
Ecrecaminrev	CATCACACCAATTTTCATACGG	<i>E. coli recA</i>
EctdkQPfor	TTTGGTGCCGGGAAAGTC	<i>E. coli tdk</i>
EctdkQPrev	CTTGTTGTCTGGTTAAAACTGG	<i>E. coli tdk</i>
<u>Nco</u> I <i>MAGP</i> NI <i>C</i>	<u>CCATGG</u> CCGGCCAGCTAGAGGAGGGGGTTCTCACC	<i>ni</i> C
<u>Nco</u> I <i>MDAP</i> NI <i>C</i>	<u>CCATGG</u> ACGCGCCAGCTAGAGGAGGGGGTTCTCACC	<i>ni</i> C
<u>Nco</u> I <i>MDGAN</i> NI <i>C</i>	<u>CCATGG</u> ACGCGGGCAGCTAGAGGAGGGGGTTCTCACC	<i>ni</i> C
<u>Nco</u> I <i>MDGP</i> NI <i>C</i>	<u>CCATGG</u> ACGCGGGCCAGCTAGAGGAGGGGGTTCTCACC	<i>ni</i> C
<u>Nco</u> I <i>NI</i> C	<u>CCATGG</u> CTAGAGGAGGGGGTTCTCACC	<i>ni</i> C
<i>tdk</i> prext2	AATAAAAATAAAGCTGAGCC	<i>tdk</i>
XntdkQPfor2	CTATCCGCTGATGCTTTGTTG	<i>tdk</i>
XntdkQPrev2	TACAATTCACAAAGCTGCTC	<i>tdk</i>

^aAdditional oligonucleotides used in this study are listed in Table 2

^bUnderlined nucleotides indicate engineered restriction sites used in cloning

^cOligonucleotide primers are specific for *X. nematophila* unless noted otherwise; Integrated DNA Technologies (Coralville, IA) synthesized the oligonucleotides used in this study.

^dItalicized nucleotides indicate *clpP*-region sequence

Conclusion

A version of *X. nematophila* Tdk protein with four extra amino acids (long-form Tdk) was not detected by immunoblot analysis of cells carrying plasmid pETLTdk even though the cells could express Tdk lacking these extra amino acids (short-form Tdk encoded on pETSTdk) under the same non-inducing growth conditions. Even with arabinose induction of transcription, long-form Tdk protein was present at a level ~13-fold lower than the short-form. The same extension reduced production of another protein, NilC, suggesting the effect of the extension is not specific to Tdk. Long-form Tdk protein was as stable as short-form Tdk protein, and its levels were not affected by the absence of Lon or ClpP proteases. Therefore the difference in long versus short-Tdk protein levels is not likely due to differential proteolysis of the two protein forms. The long-form *tdk* mRNA transcript was present at a 1.7-fold lower level than short-form *tdk* transcript and increased levels of transcript are sufficient to overcome low protein levels, supporting the hypothesis that the low level of long-form Tdk production is due in part to low long-form *tdk* transcript levels. The bacterial N-end rule [8] states that proteins containing N-terminal arginine, lysine, leucine, phenylalanine, tyrosine and tryptophan residues are particularly sensitive to Clp-dependent degradation, reducing their *in vivo* half-lives. The phenomenon reported herein is distinct from the N-end rule as none of the known destabilizing residues occur in the MDGP N-terminal

extension, nor does the extension result in protein degradation by Clp (Fig. 2D). Since the same pET-28a(+)-based expression system was used for both Tdk and NilC-trunc production, the observed phenotype may result from a combination of the MDGP extension and the pET-28a(+) expression system. These results may be relevant to researchers attempting to limit production of potentially toxic proteins from the pET-28a(+) expression plasmid under non-inducing conditions in an *E. coli* host, while still allowing protein overproduction under inducing conditions.

Methods

Bacterial strains

Strain KY895 ([9]; λ *tdk-1* IN(*rrnD-rrnE*)1, *ilv-276*) was obtained from the *E. coli* Genetic Stock Center, strain BL21 Star™ (DE3) (F-*ompT hsdS_B* (r_B⁻ m_B⁻) *gal dcm rne131* (DE3)) was obtained from Invitrogen (Carlsbad, CA) and strain BL21 (DE3) (F-*ompT hsdS_B* (r_B⁻ m_B⁻) *gal dcm* (DE3)) was obtained from Novagen (Madison, WI). To construct the *E. coli* KY895 *clpP* mutant strain (HGB871), primers clpPtetKWfor and clpPtetKWrev (Table 1) were used to amplify the tetracycline resistance gene from Tn10. The primers have 40 nt of *clpP* sequence at their 5' ends. The amplified fragment was electroporated into *E. coli* DY330 (W3110 Δ *lacU169 gal490 λ 1857 Δ (cro-bioA)*) as described elsewhere [10]. Under the conditions used, this strain allowed homologous recombination of the linear

Table 2: *X. nematophila* tdk plasmid design and resulting Tdk protein levels

Oligonucleotide name	Oligonucleotide sequence (5'→3') ^a	Resulting plasmid name	N-terminus of engineered protein ^b	Relative amount of protein expressed (% of short-form) ^c
<u>Nco</u> I Tdkshort	<u>CCATGGCTCAGCTTTATTTTAT</u>	pETSTdk	MAQLY...	100
<u>Nco</u> I Tdklong	<u>CCATGGACGGGGCCAATGGCTCAG</u>	pETLTdk	MDGPMAQLY...	0
<u>Nco</u> I MDGATdk	<u>CCATGGACGGGGCCAATGGCTCAG</u>	pETMDGATdk	MDGAMAQLY...	0
<u>Nco</u> I MDPGTdk	<u>CCATGGACCCAGGGATGGCTCAGCTTTA</u> TTTTTATTATTCTGC	pETMDPGTdk	MDPGMAQLY...	0
<u>Nco</u> I MDG(GGU)PTdk	<u>CCATGGACGGTCCAATGGCTCAG</u>	pETMDG(GGU)PTdk	MDGPMAQLY...	0
<u>Nco</u> I MAAGTdk	<u>CCATGGCCGCCGGGATGGCTCAGCTTTA</u> TTTTTATTATTCTG	pETMAAGTdk	MAAGMAQLY...	0
<u>Nco</u> I MDAGTdk	<u>CCATGGACGGCCGGGATGGCTCAGCTTTA</u> TTTTTATTATTCTG	pETMDAGTdk	MDAGMAQLY...	0.2
<u>Nco</u> I MAGPTdk	<u>CCATGGCCGGGGCCAATGGCTCAG</u>	pETMAGPTdk	MAGPMAQLY...	1
<u>Nco</u> I MGPDTdk	<u>CCATGGGGCCAGACATGGCTCAGCTTTA</u> TTTTTATTATTCTGC	pETMGPDTdk	MGPDMAQLY...	13
<u>Nco</u> I MAAAGTdk	<u>CCATGGCCGCCGCCGGGATGGCTCAGC</u> TTTATTTTATTATTCTG	pETMAAAGTdk	MAAAGMAQLY ...	15
<u>Nco</u> I MAPGDTdk	<u>CCATGGCCCCAGGGGACATGGCTCAGC</u> TTTATTTTATTATTCTG	pETMAPGDTdk	MAPGDMAQLY ...	47
<u>Nco</u> I MGDPTdk	<u>CCATGGGGGACCCAATGGCTCAGCTTTA</u> TTTTTATTATTCTGC	pETMGDPTdk	MGDPMAQLY...	77
<u>Nco</u> I MGAATdk	<u>CCATGGGGGCCGCCATGGCTCAGCTTTA</u> TTTTTATTATTCTG	pETMGAATdk	MGAAMAQLY...	80
<u>Nco</u> I MAGATdk	<u>CCATGGCCGGGGCCATGGCTCAGCTTTA</u> TTTTTATTATTCTG	pETMAGATdk	MAGAMAQLY...	88
<u>Nco</u> I MDAPTdk	<u>CCATGGACGCGCCAATGGCTCAG</u>	pETMDAPTdk	MDAPMAQLY...	93
<u>Nco</u> I MAPDGTdk	<u>CCATGGCCCCAGACGGGATGGCTCAGC</u> TTTATTTTATTATTCTG	pETMAPDGTdk	MAPDGMALY ...	103

^aUnderlined nucleotides indicate engineered restriction sites used in cloning ^bAn italicized *M* indicates an N-terminal *f*-Met residue that, based on the identity of the second amino acid residue, is predicted to be posttranslationally removed by PepM ^cAmount of Tdk protein detected by immunoblot analysis of *E. coli* KY895 carrying pTara and the pET-28a(+)-derived plasmid listed and expressed relative to the level of detected short-form Tdk and background corrected against the pET-28a(+) vector-only level.

PCR product into the DY330 chromosomal *clpP* locus, replacing the region beginning 2 bp 5' of the *clpP* start codon and ending 2 bp 3' of the *clpP* stop codon with a tetracycline resistance marker. P1 transduction was used to move the mutated locus into *E. coli* strain KY895 using standard techniques and selecting for desired transductants on tetracycline. PCR analysis was used to confirm the insertion.

Growth conditions

Cultures were grown in a tube roller at 30°C in Luria-Bertani (LB) broth [11] except for the FUDR growth assays, which were performed in semi-defined medium containing 5-fluorodeoxyuridine ([12]; FUDR obtained from Fisher Scientific, Pittsburgh, PA) at 37°C with shaking in a microplate reader (Molecular Devices, Sunnyvale, CA) as described previously [3]. LB agar (20 g l⁻¹) plates and all liquid media were supplemented when appropriate with kanamycin (kan; 20 µg ml⁻¹), chloramphenicol (cam; 20 µg ml⁻¹), isopropyl-β-D-thiogalactopyranoside (IPTG; 0.2 mM) or arabinose (0.2%). M9 medium was prepared as described elsewhere [13]. Permanent stocks of cultures

were stored at -80°C in dark-stored LB broth supplemented with 10% dimethylsulfoxide.

Production of recombinant *X. nematophila* Tdk forms in *E. coli*

DNA manipulations and transformation of *E. coli* were performed using standard protocols ([14] and product literature). To construct plasmids expressing various forms of *X. nematophila* ATCC19061 *tdk* [GenBank:AY363171] [15], the BamHItdk primer (Table 1) was combined with each primer listed in Table 2 in a PCR using either *X. nematophila* (HGB007; laboratory stock of ATCC19061) chromosomal DNA or a cloned copy of *X. nematophila* *tdk* as template DNA. Where a proline residue was desired ahead of other tested residues, an extra alanine codon was engineered between the methionine and proline codons to allow the NcoI restriction site (which includes the translational start site) to be used for cloning. The amplified products from each primer set were cloned into plasmid pET-28a(+) (Novagen, Madison, WI) using the engineered NcoI and BamHI sites. The resulting plasmids, listed in Table 2, in addition to plasmid pTara (arabinose-

inducible expression of the T7 RNA polymerase; [16]), were transformed into *E. coli* KY895, BL21 Star™ (DE3), KY895 *clpP* and BL21 (DE3). As a control, pET-28a(+) with no insert was transformed with plasmid pTara into these strain backgrounds.

Production of truncated *X. nematophila NilC* (*NilC-trunc*) forms in *E. coli*

A portion of *X. nematophila nilC* was PCR-amplified from *X. nematophila* genomic DNA using primers NcoINilC or NcoIMDGNilC and BamHINilC (Table 1), resulting in *nilC* production starting 3', at codon 22, of its predicted signal sequence-encoding region and with an additional 5' alanine codon (GCU). NcoIMDGNilC encodes an additional four codons, for the amino acid series MDGP, before the added alanine codon, to create long-form NilC. Primers NcoIMAGPNilC, NcoIMDAPNilC and NcoIMDGANilC (Table 1) were used to encode the additional amino acid series MAGP, MDAP, and MDGA, respectively, before the added alanine codon. Products from the amplification reactions were cloned into pET-28a(+) as described above. The resulting plasmids, pETnilC, pETMDGNilC, pETMAGPNilC, pETMDAPNilC and pETMDGANilC, were transformed with pTara into KY895 for protein production analysis.

Immunoblot detection of *Tdk* and *NilC*

For immunoblot detections, samples from overnight cultures of KY895 carrying pTara and the appropriate pET-28a(+)-derived vector were electrophoresed and transferred to 0.2 µm PVDF membrane (Bio-Rad) using standard protocols. Immunoblots were performed using the ECL Plus Western Blotting Kit (Amersham Biosciences, Piscataway, NJ) and a goat anti-rabbit IgG horse radish peroxidase-conjugated secondary antibody (Pierce, Rockford, IL). Anti-Tdk serum was obtained from a bleed of a New Zealand White rabbit at the University of Wisconsin Laboratory Animal Resources polyclonal antibody facility following a series of injections of purified 6× histagged *X. nematophila* Tdk [17]. Anti-Tdk and anti-*NilC* sera [5] were used at a final dilution of 1:5,000. Fluorescence was detected on a Storm860 Phosphorimager (Amersham Biosciences, Piscataway, NJ).

Primer extension mapping of short-and long-form transcripts

Total-cell RNA was isolated from KY895 carrying plasmids pTara and pETSTdk or pETLTdk following overnight growth in LB/kan/cam broth to OD₆₀₀ = 2.0. A PAGE-purified primer, tdkprext2 (Table 1), was end-labeled using T4 Polynucleotide Kinase and [^γ32-P]ATP (Perkin-Elmer Biosciences, Wellesley, MA) for 10 min at 37°C and the labeled primer used in cycle sequencing reactions with components of the *fmol*® DNA Cycle Sequencing System kit (Promega, Madison, WI) and either plasmid pETSTdk

or pETLTdk, as indicated in Figure 3. Labeled primer was also hybridized to 5 µg of total-cell RNA by dissociation at 80°C for 10 min, followed by a slow cooling to 37°C. The primer was extended by avian myeloblastosis virus reverse transcriptase enzyme (AMV-RT, Promega, Madison, WI) at 37°C and the extension products and completed sequencing reactions were resolved on a 12% SDS-polyacrylamide gel containing 8 M urea. The resulting gel was dried and visualized by exposure to a phosphor screen overnight, which was scanned on a Storm 860 phosphorimager (Amersham Biosciences, Piscataway, NJ), and the data analyzed with ImageQuant software (Amersham Biosciences, Piscataway, NJ).

Quantitative PCR to measure relative transcript levels

E. coli KY895 carrying plasmids pTara and either pET-28a(+), pETSTdk or pETLTdk was subcultured from an overnight culture and grown to OD₆₀₀~0.85 in LB broth with 0.2% glucose. Two independent cultures started from individual colonies were used for each strain. Total RNA was isolated using TRIzol (Invitrogen, Carlsbad, CA) and the RNA was DNase treated and used to make cDNA with random hexamer primers (Integrated DNA Technologies, Coralville, IA) and AMV-RT. As a control to detect DNA contamination of the DNased RNA, samples with no added AMV-RT were analyzed by PCR for amplification of *E. coli tdk* using primers EctdkQPfor and EctdkQPrev (Table 1) and, as expected, none showed product. Reactions for real-time PCR were performed in duplicate in a total volume of 25 µl with iQ™ SYBR® Green Supermix (Bio-Rad, Hercules, CA), cDNA template, appropriate primers and a three-step cycling protocol on a Bio-Rad iCycler and analyzed with Bio-Rad iCycler iQ™ software. The amount of *X. nematophila tdk* transcript was measured by amplification with XntdkQPfor2 and XntdkQPrev2 primers (Table 1). As a negative control, water was used in place of cDNA template. Cycle threshold results for each sample were adjusted according to *E. coli recA* levels (amplified with Ecrecaminf and Ecrecaminrev primers, Table 1, designed from published *E. coli recA* sequence) and then converted to arbitrary units factoring in a two-fold change in PCR product per cycle.

Pulse labeling and immunoprecipitation

Cells were grown to OD₆₀₀ = 0.3 in LB/kan/cam, washed and resuspended in M9 containing 0.4% arabinose, 1 µg ml⁻¹ thiamine, kan and cam and incubated for 1 h before being pulse labeled with 20 mCi/ml [³⁵S]-L-methionine (PerkinElmer Life Sciences, Boston, MA) for 3 min at 37°C. Unlabeled L-methionine was added to 300 µM and 50 µl samples were removed and added to 50 µl of a 125 mM Tris-Cl, 4% SDS, pH 6.8 buffer at 20 s, 2 min and 5 min after addition of unlabeled methionine. After immediate freezing in dry ice/ethanol, samples were boiled for 4 min and to each was added 1 ml of immunoprecipita-

tion buffer (50 mM Tris-Cl, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, pH7.5). The samples were centrifuged and supernatants incubated with 2 µl rabbit anti-Tdk serum with rocking for 1 h at 4°C. The samples were then incubated with Protein A immobilized on sepharose CL-4B (Sigma, St. Louis, MO) for 1 h at 4°C, with rocking. The pelleted beads were washed 3 times in immunoprecipitation buffer containing 0.1% SDS, separated on a 12% acrylamide denaturing gel and the gel was dried and exposed to a phosphor screen and analyzed as for the primer extension reactions.

Authors' contributions

SSO performed the experiments and participated in experimental design and manuscript drafting and editing. HGB participated in experimental design and manuscript drafting and editing. Both authors read and approved the final manuscript.

Acknowledgements

The authors gratefully acknowledge Charles E. Cowles for providing anti-NilC antibody. This work was supported by National Institutes of Health ROI grant GM59776, the Investigators in Pathogenesis of Infectious Disease Award from the Burroughs Wellcome Foundation and by USDA/CREES grant CRHF-0-6055 (awarded to HGB and used to support SSO in part). SSO received additional support through the National Institutes of Health Predoctoral Training Grant T32 GM07215 in Molecular Biosciences and through the National Science Foundation Graduate Teaching Fellows in K-12 Education award DUE-9979628 to the K-Through-Infinity program.

References

- Sati SP, Singh SK, Kumar N, Sharma A: **Extra terminal residues have a profound effect on the folding and solubility of a Plasmodium falciparum sexual stage-specific protein over-expressed in Escherichia coli.** *Eur J Biochem* 2002, **269**:5259-5263.
- Gottesman S: **Minimizing proteolysis in Escherichia coli: genetic solutions.** *Methods Enzymol* 1990, **185**:119-129.
- Orchard SS, Goodrich-Blair H: **Pyrimidine nucleoside salvage confers an advantage to Xenorhabdus nematophila in its host interactions.** *Appl and Environ Microbiol* in press.
- Neuhard J: **Utilization of preformed pyrimidine bases and nucleosides.** In *Metabolism of Nucleotides, Nucleosides and Nucleobases in Microorganisms* Edited by: Munch-Peterson A. London, Academic press; 1983:95-148.
- Miller JH: **Experiments in Molecular Genetics.** Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press; 1972:466.
- Cowles CE, Goodrich-Blair H: **Characterization of a lipoprotein, NilC, required by Xenorhabdus nematophila for mutualism with its nematode host.** *Mol Microbiol* 2004, **54**:464-477.
- Grunberg-Manago M: **Messenger RNA stability and its role in control of gene expression in bacteria and phages.** *Annu Rev Gen* 1999, **33**:193-227.
- Sherman F, Stewart JW, Tsunasawa S: **Methionine or not methionine at the beginning of a protein.** *BioEssays* 1985, **3**:27-31.
- Tobias JW, Shrader TE, Rocap G, Varshavsky A: **The N-end rule in bacteria.** *Science* 1991, **254**:1374-1377.
- Hiraga S, Igarashi K, Yura T: **A deoxythymidine kinase-deficient mutant of Escherichia coli. I. Isolation and some properties.** *Biochim Biophys Acta* 1967, **145**:41-51.
- Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL: **An efficient recombination system for chromosome engineering in Escherichia coli.** *Proc Natl Acad Sci USA* 2000, **97**:5978-5983.
- Summers WC, Raksin P: **A method for selection of mutations at the tdk locus in Escherichia coli.** *J Bacteriol* 1993, **175**:6049-6051.
- Sambrook J, Fritsch EF, Maniatis T: **Molecular Cloning: a Laboratory Manual.** 2nd edition. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press; 1989:1076.
- Ausubel FA, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K: **Current Protocols in Molecular Biology.** New York, Wiley and Sons; 1998.
- Orchard SS, Goodrich-Blair H: **Identification and functional characterization of a Xenorhabdus nematophila oligopeptide permease.** *Appl Env Microbiol* 2004, **70**:5621-5627.
- Wycuff DR, Matthews KS: **Generation of an AraC-araBAD promoter-regulated T7 expression system.** *Anal Biochem* 2000, **277**:67-73.
- Orchard SS: **Oligopeptide and Nucleoside Salvage by the Bacterium Xenorhabdus nematophila.** In *Bacteriology* Madison, University of Wisconsin; 2005.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

